

# Different approaches to Kovalevskaya top

Katarina Kukić\*

## Abstract

In this paper we study the equations of motion of a rigid body around a fixed point in the case of Kovalevskaya. We give interpretation of the "mysterious substitution of variables" in Sofia Kovalevskaya's paper. The present paper also connects integration procedure proposed by Golubov with the original equations obtained by S. Kovalevskaya.

**Keywords:** Kovalevskaya top, integrable case, integration procedure

## 1 Introduction

It is well known that Kovalevskaya top is one of three famous integrable cases of spinning top. This paper deals with a systematic approach to the integration of this case. Sofia Kovalevskaya integrates the problem in terms of hyperelliptic quadratures after a complicated and mysterious substitution of variables, [7]. Here we present possible explanation for this substitution and explicit formulas for the solutions obtained by adapting Golubov's method [1] to Kovalevskaya's equations from [7]. This problem was studied in detail in thesis [11].

---

\*Faculty of Transport and Traffic Engineering, University of Belgrade, e-mail: k.mijailovic@sf.bg.ac.rs (This paper got award "Rastko Stojanovic" on the First Serbian (26th YU) Congress on Theoretical and Applied Mechanics Kopaonik, Serbia, April 10-13, 2007)

The organization of this paper is as follows: Section I contains formulation of the problem and basic facts concerning Kovalevskaya's case. Section II is devoted to change of variables in the paper of Kovalevskaya. It contains Weil's interpretation of Euler's results [4], [5] [9], [10] and paraphrase of these classical results given by Jurdjevic in [6] leading to the differential equations that appear in the paper of Sofia Kovalevskaya. Section III explains procedure for integration proposed by Golubov, but adapted to equations that appear in Kovalevskaya's paper.

## 2 Formulation of the problem

A spinning top is by definition a rigid body that rotates in a constant gravitational field. The most fascinating integrable case of spinning top was discovered by Sofia Kovalevskaya in 1889 under the conditions that  $I_1 = I_2 = 2I_3, I_3 = 1, z_0 = y_0 = 0$ . With  $(I_1, I_2, I_3)$  we denote principal moments of inertia,  $(x_0, y_0, z_0)$  is center of mass,  $c = Mgx_0$  and  $(p, q, r)$  is vector of angular velocity. Finally with  $\gamma, \gamma', \gamma''$  we denote cosine of angles between  $z$  axes of fixed coordinate system and axes of coordinate system that is attached to the top and whose origin coincides with the fixed point. Then the equations of motion take the following form:

$$\begin{aligned} 2\dot{p} &= qr, & \dot{\gamma} &= r\gamma' - q\gamma'', \\ 2\dot{q} &= -pr - c\gamma'', & \dot{\gamma}' &= p\gamma'' - r\gamma, \\ \dot{r} &= c\gamma', & \dot{\gamma}'' &= q\gamma - p\gamma'. \end{aligned} \quad (1)$$

Sofia Kovalevskaya begins her investigations with solving problem for what values of  $(I_1, I_2, I_3, x_0, y_0, z_0)$  Euler-Poisson's equations admit Laurent series solutions of complex time for their general solution. Such a formulation of the problem gave one new approach to a larger class of Hamiltonian systems on Lie groups ( [6], [8], [2] ).

### 2.1 Constants of motion

System (1) has three well known independent constants of motion:

$$\begin{aligned} 2(p^2 + q^2) + r^2 &= 2c\gamma + 6l_1 - \text{energy}, \\ 2(p\gamma + q\gamma') + r\gamma'' &= 2l - \text{angular momentum in the direction of gravity}, \\ \gamma^2 + \gamma'^2 + \gamma''^2 &= 1 - \text{length of the gravity vector}. \end{aligned}$$

The extra constant of motion in the case of S. Kovalevskaya is

$$[(p + iq)^2 + c(\gamma + \nu\gamma')][(p - iq)^2 + c(\gamma - \nu\gamma')] = k^2. \quad (2)$$

Next, if we denote with

$$\begin{aligned} x_1 &= p + iq, & y_1 &= \gamma + \nu\gamma', \\ x_2 &= p - iq, & y_2 &= \gamma - \nu\gamma', \\ \xi_1 &= (p + iq)^2 + c(\gamma + \nu\gamma') = x_1^2 + cy_1, \\ \xi_2 &= (p - iq)^2 + c(\gamma - \nu\gamma') = x_2^2 + cy_2, \end{aligned}$$

then integral (2) becomes

$$\xi_1 \xi_2 = k^2. \quad (3)$$

Rewrite  $r^2, r\gamma'', \gamma''^2$  using  $x_1, x_2, \xi_1, \xi_2$ . From constants of motion and

$$\begin{aligned} p^2 + q^2 &= x_1 x_2, \\ 2c\gamma &= \xi_1 + \xi_2 - x_1^2 - x_2^2, \\ 2c\nu\gamma' &= \xi_1 - \xi_2 - x_1^2 + x_2^2, \end{aligned}$$

we get

$$\begin{aligned} r^2 &= 6l_1 - (x_1 + x_2)^2 + \xi_1 + \xi_2, \\ cr\gamma'' &= 2lc + x_1 x_2 (x_1 + x_2) - x_2 \xi_1 - x_1 \xi_2, \\ c^2 \gamma''^2 &= c^2 - k^2 - x_1^2 x_2^2 + x_2^2 \xi_1 + x_1^2 \xi_2. \end{aligned}$$

Further S. Kovalevskaya denotes with:  $E = 6l_1 - (x_1 + x_2)^2$ ,  $F = 2cl + x_1 x_2 (x_1 + x_2)$  i  $G = c^2 - k^2 - x_1^2 x_2^2$ .

Then we get

$$\begin{aligned} r^2 &= E + \xi_1 + \xi_2, \\ cr\gamma'' &= F - x_2 \xi_1 - x_1 \xi_2, \\ c^2 \gamma''^2 &= G + x_2^2 \xi_1 + x_1^2 \xi_2. \end{aligned}$$

Replacing these expressions into identity  $r^2 c^2 \gamma''^2 = (cr\gamma'')^2$  it turns into

$$(E + \xi_1 + \xi_2)(G + x_2^2 \xi_1 + x_1^2 \xi_2) - (F - x_2 \xi_1 - x_1 \xi_2)^2 = 0. \quad (4)$$

After an algebraic transformation (4) becomes

$$\begin{aligned} & [(\xi_1 + \xi_2)(x_2^2\xi_1 + x_1^2\xi_2) - (x_2\xi_1 + x_1\xi_2)^2] + [(\xi_1 + \xi_2)G + \\ & (x_2^2\xi_1 + x_1^2\xi_2)E + 2(x_2\xi_1 + x_1\xi_2)F] + EG - F^2 = 0. \end{aligned} \quad (5)$$

Using identity  $(\xi_1 + \xi_2)(x_2^2\xi_1 + x_1^2\xi_2) - (x_2\xi_1 + x_1\xi_2)^2 = \xi_1\xi_2[x_1^2 + x_2^2 - 2x_1x_2]$  from (5) we get

$$\xi_1(G + x_2^2E + 2x_2F) + \xi_2(G + x_1^2E + 2x_1F) + EG - F^2 + k^2(x_1 - x_2)^2 = 0. \quad (6)$$

Next S. Kovalevskaya denotes with:

$$\begin{aligned} R(x_1) &= x_1^2E + 2x_1F + G = x_1^2(6l_1 - (x_1 + x_2)^2) + \\ & 2x_1(2cl + x_1x_2(x_1 + x_2)) + c^2 - k^2 - x_1^2x_2^2 \\ &= [-(x_1 + x_2)^2x_1^2 + 2x_1^2x_2(x_1 + x_2) - x_1^2x_2^2] + 6l_1x_1^2 + 4clx_1 + c^2 - k^2 \\ &= -[(x_1 + x_2)x_1 - x_1x_2]^2 + 6l_1x_1^2 + 4clx_1 + c^2 - k^2 \\ &= -x_1^4 + 6l_1x_1^2 + 4clx_1 + c^2 - k^2. \end{aligned}$$

Note here that even if coefficients  $E, F, G$  depend on  $x_1$  and  $x_2$  expression  $Ex_1^2 + 2Fx_1 + G$  depends just on  $x_1$ .

Let us also introduce

$$\begin{aligned} R(x_1, x_2) &= Ex_1x_2 + F(x_1 + x_2) + G = [6l_1 - (x_1 + x_2)^2]x_1x_2 + \\ & [2cl + x_1x_2(x_1 + x_2)](x_1 + x_2) + [c^2 - k^2 - x_1^2x_2^2] \\ R_1(x_1, x_2) &= EG - F^2 = -6l_1x_1^2x_2^2 - (c^2 - k^2)(x_1 + x_2)^2 - \\ & 4lc(x_1 + x_2)x_1x_2 + 6l_1(c^2 - k^2) - 4l^2c^2. \end{aligned}$$

Then (6) becomes

$$R(x_2)\xi_1 + R(x_1)\xi_2 + EG - F^2 + k^2(x_1 - x_2)^2 = 0. \quad (7)$$

In order to express  $\xi_1$  and  $\xi_2$  as functions of  $x_1, x_2$  after short calculation and with help of identity  $R(x_1)R(x_2) - R^2(x_1, x_2) = (EG - F^2)(x_1 - x_2)^2$ , S. Kovalevskaya gets quadratic equation

$$R(x_2)\xi_1^2 + \xi_1(R_1(x_1, x_2) + k^2(x_1 - x_2)^2) + R(x_1)k^2 = 0.$$

As a next step she made the following "mysterious" substitution of variables:

$$\begin{aligned} s_1 &= \frac{R(x_1, x_2) - \sqrt{R(x_1)R(x_2)}}{2(x_1 - x_2)^2} + \frac{1}{2}l_1, \\ s_2 &= \frac{R(x_1, x_2) + \sqrt{R(x_1)R(x_2)}}{2(x_1 - x_2)^2} + \frac{1}{2}l_1. \end{aligned} \tag{8}$$

She omits any explanation concerning the above formulas and we will now try to put some light on them.

### 3 About the substitution of variables $s_{1,2} = (R(x_1, x_2) \mp \sqrt{R(x_1)R(x_2)})(x_1 - x_2)^{-2}/2 + l_1/2$

With  $P(x) = A + 4Bx + 6Cx^2 + 4Dx^3 + Ex^4$  we will denote general polynomial of degree four and with  $P_1(x, y) = A + 2B(x + y) + 3C(x^2 + y^2) + 2Dxy(x + y) + Ex^2y^2$  a particular form that satisfies  $P_1(x, x) = P(x)$ .

**Theorem 1.** The unique form that satisfies  $P_1^2(x, y) + (x - y)^2\hat{P}(x, y) = P(x)P(y)$  is  $\hat{P}(x, y) = -4B^2 + 4(AD - 3BC)(x + y) + 2(AE + 2BD - 9C^2)(x^2 + y^2) + (BE - 3CD)xy(x + y) - 4D^2x^2y^2 - (AE + 4BD - 9C^2)(x - y)^2$ .

*Proof:* By differentiation of  $F(x, y) = P(x)P(y) - P_1^2(x, y)$  we see that  $F(x, y)$  must be in the form of  $(x - y)^2\hat{P}(x, y)$ , and by further calculations one finds that  $\hat{P}(x, y)$  matches with the expression given in the statement of theorem. For more about this procedure, see [6].

Let  $P_\theta(x, y) = P_1(x, y) - \theta(x - y)^2$  denote the most general biquadratic form that satisfies  $P_\theta(x, x) = P(x)$  where  $\theta$  is an arbitrary parameter. Then,  $\Phi_\theta(x, y) = -(x - y)^2\theta^2 + 2P_1(x, y)\theta + \hat{P}(x, y)$  is the unique form that satisfies  $P_\theta^2(x, y) + (x - y)^2\Phi_\theta(x, y) = P(x)P(y)$ .

$\Phi_\theta$  can be written as  $\Phi_\theta(x, y) = a_\theta(x)y^2 + 2b_\theta(x)y + c_\theta(x) = a_\theta(y)x^2 + 2b_\theta(y)x + c_\theta(y)$  where is

$$a_\theta(x) = (2E\theta - 4D^2)x^2 + (4D\theta + 4BE - 12CD)x + AE - (\theta - 3C)^2,$$

$$b_\theta(x) = (2D\theta + 2BE - 6CD)x^2 + (\theta^2 - 9C^2 + AE + 4BD)x + 2B\theta + 2AD - 3BC,$$

$$c_\theta(x) = (AE - (\theta - 3C)^2)x^2 + (4B\theta + 4AD - 12BC)x + 2A\theta - 4B^2.$$

If with  $G_\theta(x)$  we denote discriminant  $b_\theta^2 - a_\theta c_\theta$ , then  $G_\theta(x) = p(\theta)P(x)$  where is  $p(\theta) = 2\theta(\theta - 3C)^2 + 2\theta(4BD - AE) + 4B^2E + 4AD^2 - 24BCD$ .

Polynomial  $p(\theta)$  can be linked with cubic elliptic curve

$$\Gamma = \{(\xi, \eta) : \eta^2 = 4\xi^3 - g_2\xi - g_3\},$$

where  $g_2$  and  $g_3$  are the invariants of curve  $C = \{(x, u) : u^2 = P(x)\}$  explicitly given with  $g_2 = AE - 4BD + 3C^2$  and  $g_3 = ACE + 2BDC - AD^2 - B^2E - C^3$ . If we take  $\theta = 2(\xi + C)$  and  $\eta^2 = \frac{p}{4}$  then we obtain  $\eta^2 = 4\xi^3 - g_2\xi - g_3$ .

**Theorem 2.** For each number  $\theta$ ,  $\Phi_\theta(x, y) = 0$  is a solution of one of differential equations  $\frac{dx}{\sqrt{P(x)}} \pm \frac{dy}{\sqrt{P(y)}} = 0$ , (\*). Conversely, for every solution  $y(x)$  of either equation (\*) there exists a number  $\theta$  such that  $\Phi_\theta(x, y(x)) = 0$ .

*Proof:* For the proof one can see [6],[9],[4].

In papers [9] and [10] Weil pointed out that  $\Gamma$  and  $C$  may be regarded as two components of a commutative algebraic group  $(G(C), +)$ . He describes next mappings:  $(M, N) \rightarrow M + N$  from  $C \times C$  into  $\Gamma$  and  $(M, P) \rightarrow M + P$  from  $C \times \Gamma$  into  $C$ . With each solution  $\Phi_\theta(x, y) = 0$  Weil associates two transformations from  $C$  into  $C$ , depending whether they change  $\frac{dx}{u}$  into  $\frac{dy}{v}$  or into  $-\frac{dy}{v}$ . For each point  $M = (x, u)$  on  $C$  define two points  $N_1 = (y_1, v_1)$  and  $N_2 = (y_2, v_2)$  on  $C$  by formulas:

$$y_1 = \frac{1}{a_\theta(x)}(-b_\theta(x) + 2\eta u), \quad v_1 = -\frac{1}{2\eta}(b_\theta(y_1) + a_\theta(y_1)x)$$

and

$$y_2 = \frac{1}{a_\theta(x)}(-b_\theta(x) - 2\eta u), \quad v_2 = -\frac{1}{2\eta}(b_\theta(y_2) + a_\theta(y_2)x)$$

Then, it follows that

$$\frac{dy_1}{dx} = -\frac{\frac{\partial \Phi_\theta}{\partial x}}{\frac{\partial \Phi_\theta}{\partial y}} = \frac{2\eta v_1}{2\eta u}$$

and

$$\frac{dy_2}{dx} = -\frac{\frac{\partial \Phi_\theta}{\partial x}}{\frac{\partial \Phi_\theta}{\partial y}} = -\frac{2\eta v_2}{2\eta u}.$$

Using those mappings and group structure  $G(C)$  we obtain that for value  $\theta_1 = \frac{P_1(a, b) + \sqrt{P(a)P(b)}}{(b - a)^2}$  form  $\Phi_{\theta_1}(x, y)$  is the solution of

$$\frac{dx}{\sqrt{P(x)}} - \frac{dy}{\sqrt{P(y)}} = 0 \text{ and for } \theta_2 = \frac{P_1(a, b) - \sqrt{P(a)P(b)}}{(b - a)^2} \text{ form } \Phi_{\theta_2}(x, y)$$

is the solution of  $\frac{dx}{\sqrt{P(x)}} + \frac{dy}{\sqrt{P(y)}} = 0$ . Using these results we can

now state theorem which makes clear some formulas that appeared in the Kovalevskaya's paper [7] without any explanation.

**Theorem 3.** Let  $O_1 = (\zeta_1, \eta_1)$  and  $O_2 = (\zeta_2, \eta_2)$  denote points of  $C$  related by the formulas  $O_1 = N - M$ ,  $O_2 = N + M$ . Then,

$$\frac{d\zeta_1}{\eta_1} = -\frac{dx}{u} + \frac{dy}{v}, \quad \frac{d\zeta_2}{\eta_2} = \frac{dx}{u} + \frac{dy}{v}.$$

*Proof:* For the proof one can see [6].

Sofia Kovalevskaya claims that

$$\frac{ds_1}{\sqrt{S(s_1)}} = \frac{dx_1}{\sqrt{R(x_1)}} + \frac{dx_2}{\sqrt{R(x_2)}}, \quad \frac{ds_2}{\sqrt{S(s_2)}} = -\frac{dx_1}{\sqrt{R(x_1)}} + \frac{dx_2}{\sqrt{R(x_2)}} \quad (9)$$

where is  $S(s) = 4s^3 - g_2s - g_3$  and  $g_2 = k^2 - c^2 + 3l_1^2$ ,  $g_3 = l_1(k^2 - c^2 - l_1^2) + l^2c^2$  are invariants of the curve  $C = \{(x, u) | u^2 = R(x) = -x^4 + 6l_1x^2 + 4lcx + c^2 - k^2\}$ . That is exactly application of Theorem 3, since  $s_1 = \zeta_2$  and  $s_2 = \zeta_1$ . To show this claim, we just need to notice that instead of general polynomial  $P(x)$  here we take  $R(x)$ , so  $C = l_1$ . From (8) we get

$$\begin{aligned}
s_1 &= \\
&\frac{-x_1^2x_2^2 + 3l_1(x_1^2 + x_2^2) + 2lc(x_1 + x_2) + c^2 - k^2 - 3l_1(x_1 - x_2)^2 - \sqrt{P(x_1)P(x_2)}}{2(x_1 - x_2)^2} \\
&\quad + \frac{1}{2}l_1 \\
&= \frac{P_1(x_1, x_2) - 3l_1(x_1 - x_2)^2 - \sqrt{P(x_1)P(x_2)}}{2(x_1 - x_2)^2} + \frac{1}{2}l_1 \\
&= \frac{P_1(x_1, x_2) - \sqrt{P(x_1)P(x_2)}}{2(x_1 - x_2)^2} - \frac{3l_1}{2} + \frac{l_1}{2} \\
&= \frac{\theta_2}{2} - l_1 \\
&= \frac{2(\zeta_2 + C)}{2} - l_1 \\
&= \zeta_2
\end{aligned}$$

By analogous derivation, we get  $s_2 = \zeta_1$ . Next, we will derive system of differential equations for  $s_1$  and  $s_2$ .

Let  $(x_1, \sqrt{R(x_1)})$  and  $(x_2, \sqrt{R(x_2)})$  denote points on  $C$  and  $(s_1, \sqrt{S(s_1)})$ ,  $(s_2, \sqrt{S(s_2)})$  points on  $\Gamma$ .

From  $2i\dot{x}_1 = rx_1 + c\gamma''$  we get

$$-4\left(\frac{dx_1}{dt}\right)^2 = r^2x_1^2 + 2rc\gamma''x_1 + c^2\gamma''^2.$$

Now, remember that  $r^2 = E + \xi_1 + \xi_2$ ,  $cr\gamma'' = F - x_2\xi_1 - x_1\xi_2$  and  $c^2\gamma''^2 = G + x_2^2\xi_1 + x_1^2\xi_2$ , so we get

$$-4\left(\frac{dx_1}{dt}\right)^2 = R(x_1) + \xi_1(t)(x_1(t) - x_2(t))^2.$$

By analogous derivation is

$$-4\left(\frac{dx_2}{dt}\right)^2 = R(x_2) + \xi_2(t)(x_1(t) - x_2(t))^2$$

and

$$4\frac{dx_1}{dt}\frac{dx_2}{dt} = R(x_1, x_2).$$

Next we get

$$\begin{aligned} \frac{-4}{S_1}\left(\frac{ds_1}{dt}\right)^2 &= -\frac{4}{R(x_1)}\left(\frac{dx_1}{dt}\right)^2 - \frac{4}{R(x_2)}\left(\frac{dx_2}{dt}\right)^2 - \frac{8}{\sqrt{R(x_1)R(x_2)}}\frac{dx_1}{dt}\frac{dx_2}{dt} \\ &= 2 + (x_1 - x_2)^2\left(\frac{\xi_1}{R(x_1)} + \frac{\xi_2}{R(x_2)}\right) - \frac{2R(x_1, x_2)}{\sqrt{R(x_1)R(x_2)}} \\ &= 2 - \frac{(x_1 - x_2)^2}{R(x_1)R(x_2)}(R_1(x_1, x_2) + k^2(x_1 - x_2)^2) - \frac{2R(x_1, x_2)}{\sqrt{R(x_1)R(x_2)}} \\ &= \frac{R(x_1)R(x_2) + R^2(x_1, x_2) - 2R(x_1, x_2)\sqrt{R(x_1)}\sqrt{R(x_2)} - k^2(x_1 - x_2)^4}{R(x_1)R(x_2)} \\ &= \frac{(x_1 - x_2)^4}{R(x_1)R(x_2)}\left[\left(\frac{R(x_1, x_2) - \sqrt{R(x_1)}\sqrt{R(x_2)}}{(x_1 - x_2)^2}\right)^2 - k^2\right] \end{aligned}$$

Replacing  $\frac{R(x_1, x_2) + (-1)^i\sqrt{R(x_1)}\sqrt{R(x_2)}}{2(x_1 - x_2)^2} = s_i - \frac{l_1}{2}$  and denote with  $k_1 = \frac{l_1}{2} + \frac{k}{2}$  and  $k_2 = \frac{l_1}{2} - \frac{k}{2}$  previous expression becomes

$$\begin{aligned} -\frac{1}{S_1}\left(\frac{ds_1}{dt}\right)^2 &= \frac{(x_1 - x_2)^4}{R(x_1)R(x_2)}\left(\frac{R(x_1, x_2) - \sqrt{R(x_1)}\sqrt{R(x_2)}}{2(x_1 - x_2)^2} - \frac{k}{2}\right) \\ &\quad \left(\frac{R(x_1, x_2) - \sqrt{R(x_1)}\sqrt{R(x_2)}}{2(x_1 - x_2)^2} + \frac{k}{2}\right) \\ &= \frac{(x_1 - x_2)^4}{R(x_1)R(x_2)}(s_1 - k_1)(s_1 - k_2). \end{aligned}$$

Also,  $s_2 - s_1 = \frac{\sqrt{R(x_1)}\sqrt{R(x_2)}}{(x_1 - x_2)^2}$  so finally we get

$$\frac{ds_1}{dt} = \frac{\sqrt{\Phi(s_1)}}{s_1 - s_2}$$

where  $\Phi(s_1) = -(4s_1^3 - g_2s_1 - g_3)(s_1 - k_1)(s_1 - k_2)$ .

In the same way

$$\frac{ds_2}{dt} = \frac{\sqrt{\Phi(s_2)}}{s_2 - s_1}.$$

Finally, system of equations for  $s_1$  and  $s_2$  is

$$\frac{ds_1}{\sqrt{\Phi(s_1)}} + \frac{ds_2}{\sqrt{\Phi(s_2)}} = 0, \quad \frac{s_1 ds_1}{\sqrt{\Phi(s_1)}} + \frac{s_2 ds_2}{\sqrt{\Phi(s_2)}} = dt. \quad (10)$$

## 4 Integration procedure for Kovalevskaya top

Recall from (8) that  $s_1$  and  $s_2$  are the roots of quadratic equation

$$(x_1 - x_2)^2(s - \frac{l_1}{2})^2 - R(x_1, x_2)(s - \frac{l_1}{2}) - \frac{1}{4}R_1(x_1, x_2) = 0. \quad (11)$$

Here we need to notice that Golubov in [1]  $s_1$  and  $s_2$  obtains as roots of equation

$$(x_1 - x_2)^2(s - 3l_1)^2 - 2R(x_1, x_2)(s - 3l_1) - R_1(x_1, x_2) = 0. \quad (12)$$

Since we explained possible motivation for change of variable Sofia Kovalevskaya used in her famous paper we will now adapt method proposed by Golubov in [1] on equations given by S. Kovalevskaya in [7]. More about this adaptation one can find in [11]. Denote left side of (10) with

$$Q(w, x_1, x_2) = (x_1 - x_2)^2 w^2 - R(x_1, x_2)w - \frac{1}{4}R_1(x_1, x_2). \quad (13)$$

Let us remark here that, like in [1], we can obtain system of differential equations (10) by differentiation of  $Q(w, x_1, x_2) = 0$  and substituting

$$\left(\frac{\partial Q}{\partial w}\right)^2 = R(x_1)R(x_2), \quad \left(\frac{\partial Q}{\partial x_1}\right)^2 = R(x_2)\varphi(w), \quad \left(\frac{\partial Q}{\partial x_2}\right)^2 = R(x_1)\varphi(w),$$

where is

$$\varphi(w) = 4w^3 + 6w^2l_1 + (c^2 - k^2)w - c^2l^2 = (w + \frac{3l_1}{2})(4w^2 + c^2 - k^2) - c^2l^2.$$

We can see that  $\varphi(w) = S(s)$  for  $s = w - \frac{l_1}{2}$ . In order to apply Golubov's method, next problem was to find function  $h(w)$  so that

$$Q(w, x_1, x_2)h(w) = A^2 + \varphi(w)B$$

for some expressions  $A(x_1, x_2, w)$ ,  $B(x_1, x_2, w)$ . We found that:

$$h(w) = 4(w + \frac{3l_1}{2}), \quad A = [4(w + \frac{3l_1}{2})(\frac{x_1x_2}{2} - w) + cl(x_1 + x_2)],$$

$$B = (x_1 + x_2)^2 - 4(w + \frac{3l_1}{2}).$$

If we change  $w$  into  $\tilde{s} - \frac{3l_1}{2}$ ,  $\varphi(w)$  turns into

$$\tilde{\varphi}(\tilde{s}) = \tilde{s}(4(\tilde{s} - \frac{3l_1}{2})^2 + c^2 - k^2) - c^2l^2 = 4(\tilde{s} - \tilde{e}_1)(\tilde{s} - \tilde{e}_2)(\tilde{s} - \tilde{e}_3). \quad (14)$$

Denote  $Q(w, x_1, x_2)(x_1 - x_2)^{-2}$  with  $F(w)$ . Expanding  $F(w)$  into Taylor's polynomial we get

$$\begin{aligned} F(w) &= F(u) + (w - u)F'(u) + (w - u)^2 = \\ &= \frac{1}{(x_1 - x_2)^2} \left[ 2\sqrt{u + \frac{3l_1}{2}} \left( \frac{1}{2}x_1x_2 - u \right) + \frac{lc(x_1 + x_2)}{2\sqrt{u + \frac{3l_1}{2}}} \right]^2 + \\ &+ \frac{\varphi(u)}{(x_1 - x_2)^2} \left[ \frac{(x_1 + x_2)^2}{4(u + \frac{3l_1}{2})} - 1 \right] + (w - u) \frac{2u(x_1 - x_2)^2 - R(x_1, x_2)}{(x_1 - x_2)^2} \\ &+ (w - u)^2 = 0. \end{aligned} \quad (15)$$

Now we replace  $w = \tilde{s} - \frac{3l_1}{2}$ ,  $u = z - \frac{3l_1}{2}$  and roots of (15) denote with  $\tilde{s}_1 - z$  and  $\tilde{s}_2 - z$  so

$$\begin{aligned}
(\tilde{s}_1 - z)(\tilde{s}_2 - z) &= \left[ 2\sqrt{z} \frac{\frac{1}{2}x_1x_2 - z + \frac{3l_1}{2}}{x_1 - x_2} + \frac{lc}{2\sqrt{z}} \frac{x_1 + x_2}{x_1 - x_2} \right]^2 \\
&\quad + \frac{\tilde{\varphi}(z)}{(x_1 - x_2)^2} \left[ \frac{(x_1 + x_2)^2}{4z} - 1 \right].
\end{aligned} \tag{16}$$

If we take  $z = \tilde{e}_i$ ,  $i = 1, 2, 3$  then

$$\sqrt{\tilde{e}_i} \frac{x_1x_2 - 2\tilde{e}_i + 3l_1}{x_1 - x_2} + \frac{lc}{2\sqrt{\tilde{e}_i}} \frac{x_1 + x_2}{x_1 - x_2} = \sqrt{(\tilde{s}_1 - \tilde{e}_i)(\tilde{s}_2 - \tilde{e}_i)}. \tag{17}$$

Next, denote with

$$X = \frac{x_1x_2 + 3l_1}{x_1 - x_2} = \frac{p^2 + q^2 + 3l_1}{2qi},$$

$$Y = \frac{1}{x_1 - x_2} = \frac{1}{2qi},$$

$$Z = \frac{x_1 + x_2}{x_1 - x_2} = \frac{p}{qi}$$

and

$$P_i = \sqrt{(\tilde{s}_1 - \tilde{e}_i)(\tilde{s}_2 - \tilde{e}_i)}, \quad i = 1, 2, 3.$$

Then (17) turns into:

$$\sqrt{\tilde{e}_\alpha} X - 2\tilde{e}_\alpha \sqrt{\tilde{e}_\alpha} Y + \frac{lc}{2\sqrt{\tilde{e}_\alpha}} Z = P_\alpha. \tag{18}$$

Changing  $i = 1, 2, 3$  into (18) and solving that system, we get  $X, Y, Z$ . Then we obtain

$$\begin{aligned}
q &= \frac{1}{2iY} = \frac{-1}{2i \sum_{i=1}^3 \frac{P_i \sqrt{\tilde{e}_i}}{\tilde{\varphi}'(\tilde{e}_i)}} = \frac{i}{4 \sum_{i=1}^3 \frac{P_i \sqrt{\tilde{e}_i}}{\tilde{\varphi}'(\tilde{e}_i)}}, \\
p &= \frac{Z}{2Y} = \frac{\sum_{i=1}^3 \frac{4\sqrt{\tilde{e}_j \tilde{e}_k}}{\tilde{\varphi}'(\tilde{e}_i)} P_i}{-4 \sum_{i=1}^3 \frac{P_i \sqrt{\tilde{e}_i}}{\sqrt{2} \tilde{\varphi}'(\tilde{e}_i)}} = - \frac{\sum_{\alpha=1}^3 \frac{\sqrt{\tilde{e}_j \tilde{e}_k}}{\tilde{\varphi}'(\tilde{e}_i)} P_i}{\sum_{i=1}^3 \frac{P_i \sqrt{\tilde{e}_i}}{\tilde{\varphi}'(\tilde{e}_i)}},
\end{aligned}$$

$$r = \frac{2\dot{p}}{q} = \frac{\sum_{i=1}^3 \frac{\sqrt{\tilde{e}_i} P_{jk}}{\tilde{\varphi}'(\tilde{e}_i)}}{\iota(\sum_{i=1}^3 \frac{\sqrt{\tilde{e}_i} P_i}{\tilde{\varphi}'(\tilde{e}_i)}}$$

where  $(i, j, k)$  is cyclic permutation of  $(1, 2, 3)$ . With  $P_{ij}$  we denoted

$$P_{ij} = P_i P_j \left[ \frac{\frac{d\tilde{s}_1}{dt}}{(\tilde{s}_1 - \tilde{e}_i)(\tilde{s}_1 - \tilde{e}_j)} + \frac{\frac{d\tilde{s}_2}{dt}}{(\tilde{s}_2 - \tilde{e}_i)(\tilde{s}_2 - \tilde{e}_j)} \right]$$

for  $e_4 = k_1 + l_1$ ,  $e_5 = k_2 + l_1$ ,  $i, j = 1, 2, 3, 4, 5$ .

Next we obtain  $\gamma''$  from equations  $2\dot{p} = qr$ ,  $2\dot{q} = -pr - c\gamma''$ , so

$$qc\gamma'' = -\frac{d(p^2 + q^2)}{dt} = -\frac{d(\frac{X}{Y})}{dt}$$

and we get

$$\gamma'' = \frac{1}{\iota c} \frac{\sum_{i=1}^3 \frac{\sqrt{\tilde{e}_i \tilde{e}_j}}{\tilde{\varphi}'(\tilde{e}_k)} P_{ij}}{\sum_{i=1}^3 \frac{\sqrt{\tilde{e}_i}}{\tilde{\varphi}'(\tilde{e}_i)} P_i}.$$

From constant of energy and  $2(p^2 + q^2) = \frac{2X}{Y} - 6l_1$  we can express

$$2c\gamma = \frac{2X}{Y} + r^2 - 12l_1 = 4 \frac{\sum_{i=1}^3 \frac{\sqrt{\tilde{e}_i(\tilde{e}_j + \tilde{e}_k)} P_i}{\tilde{\varphi}'(\tilde{e}_i)}}{\sum_{i=1}^3 \frac{P_i \sqrt{\tilde{e}_i}}{\tilde{\varphi}'(\tilde{e}_i)}} - \left( \frac{\sum_{i=1}^3 L_i P_{jk}}{\sum_{i=1}^3 L_i P_i} \right)^2 - 12l_1,$$

where with  $L_i$  we denoted  $\iota(\tilde{e}_j - \tilde{e}_k)\sqrt{\tilde{e}_i}$ . Finally, we obtain  $\gamma'$  from equation  $\dot{r} = c\gamma'$ . Then,

$$\begin{aligned} & -c\gamma' (L_1 P_1 + L_2 P_2 + L_3 P_3)^2 \\ & = \sum_{i=1}^3 (\tilde{e}_j - \tilde{e}_k)^2 \tilde{e}_i P_{i4} P_{i5} \\ & + \sum_{i=1}^3 (\tilde{e}_i - \tilde{e}_j)(\tilde{e}_k - \tilde{e}_i) \sqrt{\tilde{e}_i \tilde{e}_j} (P_{j4} P_{k5} + P_{j5} P_{k4} + (\tilde{e}_j - \tilde{e}_k)^2 P_i). \end{aligned}$$

In paper [7] S. Kovalevskaya obtained formulas for  $p, q, r, \gamma, \gamma', \gamma''$  using Weierstrass  $\wp$  and  $\sigma$  function after long and not easy calculations. With  $e_1, e_2, e_3$  she denotes the roots of  $\mathcal{S}(s)$  and uses change of variable  $w = s -$

$\frac{l_1}{2}$ , so  $\tilde{e}_i = e_i + l_1$ . With these remarks it is easy to check that expressions we got here are the same as those obtained by S. Kovalevskaya. She did, also, expressed  $P_i, P_{ij}$  for  $i, j \in \{1, 2, 3, 4, 5\}$  through  $\vartheta$  functions. More about these formulas and notation she used, one can see in [3], [11].

## 5 Conclusions

Even after more than hundred years after publishing, Kovalevskaya's paper [7] is still quite a challenge. Up to now, her method has been neither understood, nor improved. This paper offered simple alternative to Kovalevskaya's formulas without mentioning properties of  $\wp$  and  $\sigma$  functions. New techniques are still developing. Some geometric approaches to this problem were proposed in [8], [2] and that will be the field of our future research.

## References

- [1] V. V. Golubev, *Lectures on the Integration of Equations of Motion of a Heavy Rigid Body Around a Fixed Point*, Gostekhizdat, 1953.
- [2] Mark Adler, Pierre van Moerbeke, Pol Vanhaecke, *Algebraic Integrability, Painlevé Geometry and Lie Algebras*, Springer Verlag, Berlin-Heidelberg, 2004.
- [3] H. F. Baker, *Abelian functions, Abel's theorem and the Allied Theory of Theta Functions*, Cambridge University Press, 1995.
- [4] L. Euler, *Evolutio generalior formularum comparationi curvarum inservientium*, Opera Omnia, Series 1<sup>a</sup>, vol. 20, pp. 318-356, 1754.
- [5] L. Euler, *De integratione aequationis differentialis  $\frac{mdx}{\sqrt{(1-x^4)}} = \frac{ndy}{\sqrt{(1-y^4)}}$* , Opera Omnia, Series 1<sup>a</sup>, vol. 20, pp. 58-79, 1754.
- [6] V. Jurdjevic, *Integrable Hamiltonian systems on Lie groups: Kowalewski type*, Annals of Mathematics, 150, p.605-644, 1999.

- [7] S. Kowalevski, Sur le problème de la rotation d'un corps solide autour d'un point fixe, *Acta Mathematica*, 12, p.177-232., 1889
- [8] A. Lesfari, Abelian surfaces and Kowalewski's top, *Annales scientifiques de l'É.N.S. 4<sup>e</sup> série*, tome 21, n<sup>o</sup>2, (1988), p. 193-223.
- [9] A. Weil, Euler and the Jacobians of elliptic curves, *Arithmetic and Geometry*, Vol. I, *Progr. Math.* 35, Birkhäuser, Boston, Mass, 353-359, 1983.
- [10] A. Weil, Remarques sur un mémoire d'Hermite, *Arch. Math.* 5, p.197-202, 1954.
- [11] K. Kukić, Klasični i savremeni pristupi zadatku Kovaljevske, Magistrski rad, Matematički fakultet, Beograd 2007.

Submitted on October 2007.

## **Razni pristupi čigri Kovaljevske**

Proučavaju se jednačine kretanja krutog tela oko nepomične tačke u slučaju Kovaljevske. Daje se interpretacija "tajanstvene smene promenljivih" u radu Sofije Kovaljevske. Ovde predloženi rad takodje povezuje integracioni postupak Golubova sa originalnim jednačinama dobijenim Sofijom Kovaljevskom.